



M-PA Agar Base

RDM-MPA-01

Principle

Levin and Cabelli (1972) described M-PA Agar as a selective membrane filter medium for *Pseudomonas aeruginosa*. The media is composed of L-lysine, yeast extract, xylose, sucrose, lactose, sodium chloride, ferric ammonium citrate, sodium thiosulphate, phenol red and agar. L-lysine is source of amino acid. Yeast extract provide nitrogen, carbon, vitamins and other necessary nutrients or growth factors. Xylose, sucrose and lactose are fermentable carbohydrates. Sodium chloride maintain osmotic balance. Ferric ammonium citrate and provides essential ions. Sodium thiosulphate is source of sulphate. Phenol red is pH indicator, which turns yellow under acidic conditions due to fermentation of the carbohydrates. Agar is solidifying agent. The selectivity of media can be increase by adding antibiotics like Sulphapyridine, Kanamycin, Nalidixic acid and Amphotericin B. The sulfonamides are synthetic bacteriostatic antibiotics with a wide spectrum against most gram-positive and many gram-negative organisms. Kanamycin inhibits protein synthesis in gram-positive organisms. Nalidixic acid blocks replication of susceptible gram-negative bacteria. Amphotericin inhibits fungal species.

Use: Recommended for the detection and isolation of *Pseudomonas aeruginosa* by membrane filter technique.

Contents*

Ingredients

	Gram/Litre
L-Lysine	5.000
Yeast Extract	2.000
Xylose	2.500
Sucrose	1.250
Lactose	1.250
Sodium Chloride	5.000
Ferric Ammonium Citrate	0.800
Sodium Thiosulphate	6.800
Phenol Red	0.080
Agar	15.000
pH at 25°C	7.1 ±0.2

* Formula adjusted for optimum performance and parameters

Directions: Dissolve 40.00 grams in 1000 ml distilled water, boil to dissolve the medium completely and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 min, cool it to 42-45 °C and add 5.0 ml of sterile antibiotic solution (Sulphapyridine (176 mg) Kanamycin (8.50 mg), Nalidixic acid (37 mg) and Amphotericin B (5.0 mg)). Mix well and distribute aseptically in sterile petri plates. Ensure complete solidification and inoculate test sample aseptically.

Specimens types analyzed

Pharmaceutical samples, clinical and non-clinical samples etc.

Precautions to be taken

These microbial media are intended for the in-vitro use only. All the handling, experiments, storage, and discarding should be performed with the help of skilled and knowledgeable technicians and as per the established guidelines. The material should be disposed only after proper sterilization by autoclaving. Please go through the MSDS of the media to avoid any accidents or in emergency.

Performance and Evaluation

The expected performance of the medium is liable to use as per the direction on the label when stored at optimum conditions and within expiry date.

Quality Control

Appearance	Light pink beige colored free flowing, homogeneous powder
Reaction of 4.00 % solution	7.1 ±0.2 at 25 °C
pH	6.90- 7.30
Gelling	Firm comparable with 1.5% agar gel
Color and clarity of ready medium	Orange red colored slightly opalescent gel
Growth Promotion properties	Best at ≤ 100 CFU at 32-37 °C for 18-72 h
Indicative properties	Optimum at ≤ 100 CFU at 32-37 °C for 18-48 h
Negative control	Performed using sterile distilled water

Different Microbial Response

Organism	ATCC	Inoculum	Growth	Incubation
<i>Pseudomonas aeruginosa</i>	27853	50-100	Luxurious	40-42°C, 48-72 h
<i>Escherichia coli</i>	8739	50-100	Inhibited	33-37 °C, 18-48 h

Storage and Shelf Life

Hygroscopic; keep container tightly closed. Store in cool dry place.

Disposal: To avoid the contamination or propagation of any hazardous microbes the used, unusable or modified preparation of this product must be disposed after autoclaving after completion of task.

Reference

1. Atlas, R. M. (2005). *Handbook of media for environmental microbiology*. CRC press.
2. Levin, M. A., and V. J. Cabelli. (1972). *Membrane filter technique for enumeration of Pseudomonas aeruginosa*. Appl. Microbiol. 24:864-870.

Disclaimer

The information contained in the technical data sheet is to the best of our knowledge is accurate and true based on the research and development work carried out by **ReadyMED**®, Chaitanya Agro Biotech, Malkapur, Maharashtra. The products are neither intended for any therapeutic use for animal or human nor for any other *in-vivo* applications. The **ReadyMED**® products are only meant to be used for the laboratory, diagnostic, research, or further manufacturing purpose only. These technical outcomes should not be considered as the warranty of any kind expressed or implied, and no liability is accepted for infringement of any patent.

CHAITANYA AGRO BIOTECH PVT. LTD. An ISO 11134:2014, ISO 13485:2016, ISO 9001:2015 CE, CIN NO.: U24210MH1995PTC095220S,
S. No. 120/2, Laxmi Nagar, Umbarnala Road, Malkapur-443101, Dist.: Buldana (M.S.) India. Customer Care +91-8669083859
rdmsales@chaitanyagroupindia.com, mkt.cabt@chaitanyagroupindia.com, www.chaitanyagroupindia.com