



Sabouraud Dextrose Agar (Harmonized)

RDM-H-SDA-01

Principle

Sabouraud dextrose agar is used for cultivation of yeasts, molds and aciduric bacteria. It is also recommended by USP, IP, JP and EP as a medium for microbial limit testing of pharmaceutical products and raw material used in pharmaceutical industries. The medium is prepared in accordance with the harmonized principles of USP/EP/IP/JP. Medium is consisting of meat and casein peptone (1:1), dextrose monohydrate and agar. The meat and casein peptone provide carbonaceous, nitrogenous compounds, long chain amino acids, vitamins and other essential growth nutrients in addition to that the dextrose monohydrate serves as energy source. The high concentration of dextrose and low pH of medium favor the growth of yeasts and molds and inhibit other contaminating bacteria from pharmaceutical and clinical specimens. Agar is used as solidifying agent.

Use: Recommended for the cultivation of yeasts, molds and aciduric bacteria from pharmaceutical products in accordance with the microbial limit testing by harmonized principles of USP/EP/BP/IP.

Contents*

Ingredients

	Gram/Litre
Meat and Casein Peptone	10.000
Dextrose Monohydrate	40.000
Agar	15.000
pH at 25°C	5.6 ±0.2

* Formula adjusted for optimum performance and parameters

Directions: Dissolve 65.0 grams in 1000 ml distilled water. Sterilize by autoclaving at 15 lbs. pressure (121°C) for 15 min, cool it to 42-45 °C and distribute aseptically in to the petri plates and allow to solidify. Ensure complete solidification and inoculate test sample aseptically.

As per IP and EP the media can be fortified with antibiotic like 0.1g of benzylpenicillin sodium and 0.1g of tetracycline per liter as sterile solution or alternatively 50 mg of chloramphenicol per liter of medium before sterilization to achieve more selectivity.

Specimens types analyzed

Pharmaceutical samples, clinical samples, *etc.*

Precautions to be taken

These microbial media are intended for the in-vitro use only. All the handling, experiments, storage, and discarding should be performed with the help of skilled and knowledgeable technicians and as per the established guidelines. The material should be disposed only after proper sterilization by autoclaving. Please go through the MSDS of the media to avoid any accidents or in emergency.

Performance and Evaluation

The expected performance of the medium is liable to use as per the direction on the label when stored at optimum conditions and within expiry date.

Quality Control

Appearance	Light beige colored free flowing, homogeneous powder
Reaction of 6.5% solution	5.6 ±0.2 at 25 °C
pH	5.40- 5.80 at 25 °C
Gelling	Firm comparable with 1.5% agar gel
Color and clarity of ready medium	Light to medium amber, very slightly opalescent without significant

	precipitate
Growth Promotion properties	Best at ≤ 100 CFU at 32-37 °C for 18-72 h
Indicative properties	Optimum at ≤ 100 CFU at 32-37 °C for 18-48 h
Negative control	Performed using sterile distilled water

Different Microbial Response

Organism	ATCC	Inoculum CFU	Observed CFU	Recovery (%)	Incubation temp. and period
<i>Candida albicans</i>	10231	50-100	Luxurious	70-75%	30-37°C, 3-5 days
<i>Aspergillus brasiliensis</i>	16404	50-100	Luxurious	70-75%	30-37°C, 3-5 days
<i>Saccharomyces cerevisiae</i>	9736	50-100	Luxurious	70-75%	30-37°C, 3-5 days

Storage and Shelf Life

Hygroscopic; keep container tightly closed. Store in cool dry place.

Disposal: To avoid the contamination or propagation of any hazardous microbes the used, unusable or modified preparation of this product must be disposed after autoclaving after completion of task.

Reference

1. Atlas, R. M. (2005). *Handbook of media for environmental microbiology*. CRC press.
2. *Difco Manual* (1998). 11th Edition. Difco Laboratories., Division of Becton Dickinson and Company, Sparks, Maryland, USA.
3. *European Pharmacopoeia*, (2011), European Dept. for the quality of Medicines.
4. *Indian Pharmacopoeia*, (2018), Govt. of India, the Controller of Publication, New Delhi.
5. Rand, M. C., Arnold E. Greenberg, and Michael J. Taras, (1976), *Standard methods for the examination of water and wastewater*. Prepared and published jointly by American Public Health Association, American Water Works Association, and Water Pollution Control Federation.
6. *The Japanese Pharmacopoeia*, 17th Ed. (2016), The Ministry of Health, Labour And Welfare
7. *The United States Pharmacopoeia*, (2014), The United States Pharmacopeial Convention. 12601 Twinbrook Parkway, Rockville, MD 20852.

Disclaimer

The information contained in the technical data sheet is to the best of our knowledge is accurate and true based on the research and development work carried out by **ReadyMED**[®], Chaitanya Agro Biotech, Malkapur, Maharashtra. The products are neither intended for any therapeutic use for animal or human nor for any other *in-vivo* applications. The **ReadyMED**[®] products are only meant to be used for the laboratory, diagnostic, research, or further manufacturing purpose only. These technical outcomes should not be considered as the warranty of any kind expressed or implied, and no liability is accepted for infringement of any patent.

CHAITANYA AGRO BIOTECH PVT. LTD. An ISO 11134:2014, ISO 13485:2016, ISO 9001:2015 CE, CIN NO.: U24210MH1995PTC095220S,
S. No. 120/2, Laxmi Nagar, Umbarnala Road, Malkapur-443101, Dist.: Buldana (M.S.) India. Customer Care +91- 8669083859
rdmsales@chaitanyagroupindia.com, mktcabt@chaitanyagroupindia.com, www.chaitanyagroupindia.com