



Tergitol 7 Agar Base

RDM-T7A-01

Principle

Tergitol 7 agar base is composed of proteose peptone, yeast extract, lactose, tergitol 7, bromothymol blue and agar. Proteose peptone provides nitrogen and amino acids. Yeast extracts serve nitrogen, carbon, and vitamin. Lactose is fermentable carbohydrate. Tergitol 7 is selective agents ease in the recovery of injured coliforms. Brom thymol bule is pH indicators. Lactose fermenting microorganism like *Escherichia coli* form yellow color colonies with yellow zone, while *Enterobacter* and *Klebsiella* colonies are greenish-yellow and lactose non-fermenting organisms, such as *Salmonella* and *Shigella*, produce colonies surrounded by blue zones. For more differentiation triphenyltetrazolium chloride is added to the medium, it serves as an indicator of lactose fermenting and non-fermenting bacterial growth. Triphenyltetrazolium chloride is rapidly reduced to insoluble red formazan by most lactose-fermenting organisms except *Escherichia coli*, *Enterobacter* and *Klebsiella* species. While other lactose fermenters, like coliforms, produce greenish-yellow colonies with yellow zones. The lactose non-fermenters produce red colonies surrounded by blue zones. Agar is a solidifying agent. The media can be fortified with Penicillin G (1.0 µg/ml), aseptically added to the medium after autoclaving, prevents growth of gram-positive cocci. The fecal coliform bacteria are identified as yellow colonies with deep yellow halo after incubation at 44°C.

Use: Recommended for recovering injured coliforms from treated water.

Contents*

Ingredients	Gram/Litre
Proteose Peptone	5.000
Yeast Extract	3.000
Lactose	10.000
Tergitol 7	0.100
Bromothymol Blue	0.025
Agar	15.000
pH at 25°C	6.9 ±0.2

* Formula adjusted for optimum performance and parameters

Directions: Dissolve 33.00 grams in 1000 ml distilled water, boil to dissolve the medium completely and sterilize by autoclaving at 15 lbs. pressure (121°C) for 15 min, cool it to 42-45 °C and distribute aseptically in petri plates. Ensure complete solidification and inoculate test sample aseptically.

Note: For more differentiation, add 4.0 ml of filter-sterilized 1% Triphenyltetrazolium chloride solution.

Specimens types analyzed

Water samples etc.

Precautions to be taken.

These microbial media are intended for the in-vitro use only. All the handling, experiments, storage, and discarding should be performed with the help of skilled and knowledgeable technicians and as per the established guidelines. The material should be disposed only after proper sterilization by autoclaving. Please go through the MSDS of the media to avoid any accidents or in emergency.

Performance and Evaluation

The expected performance of the medium is liable to use as per the direction on the label when stored at optimum conditions and within expiry date.

Quality Control

Appearance	Beige colored free flowing, homogeneous powder
Reaction of 3.3% solution	6.9 ±0.2 at 25 °C
pH	6.70- 7.10
Gelling	Firm comparable with 1.5% agar gel
Color and clarity of ready medium	Green colored slightly opalescent gel
Growth Promotion properties	Best at ≤ 100 CFU at 32-37 °C for 18-72 h
Indicative properties	Optimum at ≤ 100 CFU at 32-37 °C for 18-48 h
Negative control	Performed using sterile distilled water

Different Microbial Response

Organism	Inoculum	Growth	Acid production	Incubation Temperature	Incubation period
<i>Escherichia coli</i> (ATCC 8739)	50-100	Luxurious	Yellow colony positive	33-37 °C	18-48 h
<i>Salmonella typhimurium</i> (ATCC 14028)	50-100	Luxurious	Blue colony negative	33-37 °C	18-48 h

Storage and Shelf Life

Hygroscopic; keep container tightly closed. Store in cool dry place.

Disposal: To avoid the contamination or propagation of any hazardous microbes the used, unusable or modified preparation of this product must be disposed after autoclaving after completion of task.

Reference

1. Atlas, R. M. (2005). *Handbook of media for environmental microbiology*. CRC press.
2. *Difco Manual* (1998). 11th Edition. Difco Laboratories., Division of Becton Dickinson and Company, Sparks, Maryland, USA.

Disclaimer

The information contained in the technical data sheet is to the best of our knowledge is accurate and true based on the research and development work carried out by **ReadyMED**[®], Chaitanya Agro Biotech, Malkapur, Maharashtra. The products are neither intended for any therapeutic use for animal or human nor for any other *in-vivo* applications. The **ReadyMED**[®] products are only meant to be used for the laboratory, diagnostic, research, or further manufacturing purpose only. These technical outcomes should not be considered as the warranty of any kind expressed or implied, and no liability is accepted for infringement of any patent.

CHAITANYA AGRO BIOTECH PVT. LTD. An ISO 11134:2014, ISO 13485:2016, ISO 9001:2015 CE, CIN NO.: U24210MH1995PTC095220S,
S. No. 120/2, Laxmi Nagar, Umbarnala Road, Malkapur-443101, Dist.: Buldana (M.S.) India. Customer Care +91-8669083859
rdmsales@chaitanyagroupindia.com, mkt.cabt@chaitanyagroupindia.com, www.chaitanyagroupindia.com